

Medical Policy

Healthcare Services Department

Policy Name	Policy Number	Scope
Octreotide Agents (Sandostatin®, Sandostatin LAR®)	MP-RX-FP-65-23	<input checked="" type="checkbox"/> MMM MA <input checked="" type="checkbox"/> MMM Multihealth

Service Category

- | | |
|--|---|
| <input type="checkbox"/> Anesthesia | <input type="checkbox"/> Medicine Services and Procedures |
| <input type="checkbox"/> Surgery | <input type="checkbox"/> Evaluation and Management Services |
| <input type="checkbox"/> Radiology Procedures | <input type="checkbox"/> DME/Prosthetics or Supplies |
| <input type="checkbox"/> Pathology and Laboratory Procedures | <input checked="" type="checkbox"/> Part B DRUG |

Service Description

This document addresses the use of **Octreotide Agents (Sandostatin®, Sandostatin LAR®)**, a drug approved by the Food and Drug Administration (FDA) for the treatment of Acromegaly, Carcinoid Tumors and Vasoactive Intestinal Peptide Tumors (VIPomas).

Background Information

Octreotide acetate agents exerts pharmacologic actions similar to the natural hormone somatostatin, but is a more potent inhibitor of GH, glucagon, and insulin than somatostatin. Like somatostatin, it also suppresses luteinizing hormone (LH) response to gonadotropin-releasing hormone (GnRH), decreases splanchnic blood flow, and inhibits release of serotonin, gastrin, vasoactive intestinal peptide, secretin, motilin, and pancreatic polypeptide.

Octreotide acetate agents are approved by the FDA to treat acromegaly, to reduce flushing episodes and watery diarrhea caused by cancerous tumors (carcinoid syndrome) and tumors called vasoactive intestinal peptide tumors (VIPomas).

Acromegaly is a rare condition that occurs if a tumor causes excess growth hormone secretion, that in turn increases IGF-1 levels. The increase in the hormones causes the hands, feet, lips, nose, and tongue to become larger, bone changes, headaches, joint aches, and vision problems. Complications may develop such as type 2 diabetes, high blood pressure, heart disease, sleep apnea, and arthritis. Estimates are there are 3,000 new cases of acromegaly per year with a prevalence of about 25,000 patients in the US. Treatment includes surgery, radiation, and medications. Medications are used if surgery is impractical or not successful. Medications for acromegaly include somatostatin analogs, growth hormone receptor antagonist, and dopamine agonist. Dopamine agonist (e.g., cabergoline) has a limited role in the treatment of acromegaly for those with mild disease.

Sandostatin® and Sandostatin LAR® are used for the symptomatic treatment of severe diarrhea in patients with carcinoid tumors and vasoactive intestinal peptide-secreting tumors (VIPomas). Their effect on tumor size, growth rate, and the development of metastases has not been determined.

Octreotide acetate agents may cause cardiac function abnormalities, including an increased risk of atrioventricular blocks, bradycardia, arrhythmias, and conduction issues, warranting cardiac monitoring, especially during intravenous administration, and possible dosage adjustments of cardiac medications. It can

Medical Policy

Healthcare Services Department

Policy Name	Policy Number	Scope
Octreotide Agents (Sandostatin®, Sandostatin LAR®)	MP-RX-FP-65-23	<input checked="" type="checkbox"/> MMM MA <input checked="" type="checkbox"/> MMM Multihealth

also lead to cholelithiasis, requiring periodic monitoring and discontinuation if complications arise. Glucose metabolism disturbances, such as hypoglycemia or hyperglycemia, may occur, necessitating glucose monitoring and adjustments in anti-diabetic treatments. Additionally, hypothyroidism may develop, requiring periodic thyroid function monitoring. Patients may experience steatorrhea and malabsorption of dietary fats, requiring evaluation for pancreatic exocrine insufficiency if symptoms worsen.

The safety and/or efficacy of octreotide acetate have not been established for treating the following conditions. The peer-reviewed published medical literature consists of case reports, small case series, RCTs of small sample sizes, and non-randomized or uncontrolled trials which preclude drawing reliable conclusions on the safety and net health benefit of octreotide acetate for other conditions, including but not limited to:

1. AIDs-related diarrhea (Panel 2018);
2. Chyle fistula management following neck dissection surgery (Swanson, 2015);
3. Chylothorax in adults (Fujita, 2014; Ismail, 2015) and neonates (Das and Shah, 2010; Testoni, 2015);
4. Graves' ophthalmopathy (thyroid eye disease) (Stan, 2006);
5. Hypothalamic obesity (insulin hypersecretion) (Lustig, 2003; Michalsky, 2012);
6. Other carcinomas, such as:
 - Advanced, metastatic breast cancer (Bajetta, 2002; Chapman, 2015);
 - Hepatocellular cancer (Jia, 2010);
 - Prostate cancer (including castration-resistant) (Friedlander, 2012);
7. Other GI tract conditions, such as:
 - Bleeding from vascular malformations (such as, angiodysplasias, angioectasias, or/GI tract AVM (Brown, 2010; Junquera, 2007; Loyaga-Rendon, 2015, Szilagyi and Ghali, 2006);
 - Gastroparesis (Edmunds, 1998);
 - Non-variceal upper GI bleeding (Archimandritis, 2000);
 - Pancreatitis (Xu, 2013);
 - Short bowel syndrome (Nehra, 2001);
 - Small intestinal dysmotility associated with systemic sclerosis (scleroderma) (Nikou, 2007; Perlemuter, 1999; Soudah, 1991; Verne, 1995); and
8. Polycystic kidney or liver disease (Caroli, 2013; Hogan, 2010; Ruggerenti, 2005).

Approved Indications

Table 1 includes the FDA-approved indications for the somatostatin analogs.

Other Uses

None.

Medical Policy

Healthcare Services Department

Policy Name	Policy Number	Scope
Octreotide Agents (Sandostatin®, Sandostatin LAR®)	MP-RX-FP-65-23	<input checked="" type="checkbox"/> MMM MA <input checked="" type="checkbox"/> MMM Multihealth

Table 1: Somatostatin Analogs

Product	Indications
Sandostatin® (octreotide acetate injection)	<ul style="list-style-type: none"> Indicated for the treatment of: <ul style="list-style-type: none"> Patients with acromegaly who have had inadequate response to or cannot be treated with surgical resection, pituitary irradiation, and bromocriptine mesylate at maximally tolerated doses. Carcinoid tumors Vasoactive Intestinal Peptide Tumors (VIPomas)
Sandostatin LAR® depot (octreotide acetate injection)	<ul style="list-style-type: none"> Indicated for the treatment of patients who have responded to and tolerated Sandostatin Injection subcutaneous for: <ul style="list-style-type: none"> Acromegaly Severe diarrhea/flushing episodes associated with metastatic carcinoid tumors Profuse watery diarrhea associated with Vasoactive Intestinal Peptide (VIP) secreting tumors

Medical Policy

Healthcare Services Department

Policy Name	Policy Number	Scope
Octreotide Agents (Sandostatin®, Sandostatin LAR®)	MP-RX-FP-65-23	<input checked="" type="checkbox"/> MMM MA <input checked="" type="checkbox"/> MMM Multihealth

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Benefit coverage for health services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

HCPCS	Description
J2353	Injection, octreotide, depot form for intramuscular injection, 1 mg [Sandostatin LAR®]
J2354	Injection, octreotide, nondepot form for subcutaneous or intravenous injection, 25 mcg [Sandostatin®]
J3490	Unclassified drugs [Bynfezia Pen]
C9399	Unclassified drugs or biologicals [Bynfezia Pen]

ICD-10	Description
C18.0-C18.9	Malignant neoplasm of colon [associated bowel obstruction]
C25.0-C25.9	Malignant neoplasm of pancreas [related VIPoma syndrome]
C37	Malignant neoplasm of thymus
C48.1-C48.8	Malignant neoplasm of peritoneum [associated bowel obstruction]
C57.00-C57.4	Malignant neoplasm of other and unspecified female genital organs [associated bowel obstruction]
C70.0-C70.9	Malignant neoplasm of meninges
C74.00-C74.92	Malignant neoplasm of adrenal gland
C75.1	Malignant neoplasm of pituitary gland
C7A.00-C7A.8	Malignant neuroendocrine tumors (carcinoid tumors)
C7B.00-C7B.09	Secondary carcinoid tumors
C7B.8	Other secondary neuroendocrine tumors
D01.7	Carcinoma in situ of other specified digestive organs [pancreas]
D13.7	Benign neoplasm of endocrine pancreas
D15.0	Benign neoplasm of thymus
D32.0-D32.9	Benign neoplasm of meninges
D35.2	Benign neoplasm of pituitary gland
D3A.010-D3A.8	Benign neuroendocrine tumors
E05.80-E05.81	Other thyrotoxicosis
E22.0	Acromegaly and pituitary gigantism
E31.20-E31.23	Multiple endocrine neoplasia [MEN] syndrome

Medical Policy

Healthcare Services Department

Policy Name	Policy Number	Scope
Octreotide Agents (Sandostatin®, Sandostatin LAR®)	MP-RX-FP-65-23	<input checked="" type="checkbox"/> MMM MA <input checked="" type="checkbox"/> MMM Multihealth

E34.00-E34.09	Carcinoid syndrome
H47.49	Disorders of optic chiasm in (due to) other disorder
I85.01	Esophageal varices with bleeding
I85.11	Secondary esophageal varices with bleeding
K56.690-K56.699	Other intestinal obstruction
K59.1	Functional diarrhea
K70.0-K75.9	Disease of liver [related bleeding esophageal varices
R19.7	Diarrhea, unspecified
Z85.841	Personal history of malignant neoplasm of brain
Z85.845	Personal history of malignant neoplasm of other parts of nervous tissue

Medical Policy

Healthcare Services Department

Policy Name	Policy Number	Scope
Octreotide Agents (Sandostatin®, Sandostatin LAR®)	MP-RX-FP-65-23	<input checked="" type="checkbox"/> MMM MA <input checked="" type="checkbox"/> MMM Multihealth

Medical Necessity Guidelines

When a drug is being reviewed for coverage under a member's medical benefit plan or is otherwise subject to clinical review (including prior authorization), the following criteria will be used to determine whether the drug meets any applicable medical necessity requirements for the intended/prescribed purpose.

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

Clinical Criteria:

B vs D Criteria: All drugs included in this PA are subject to B vs D evaluation. Medication must be furnished "incident to" physician service provided and usually not self-administered to be covered by Medicare and to be eligible to be evaluated through part B. If not, medication must be evaluated through part D.

Octreotide (Bynfezia®, Sandostatin® or Sandostatin LAR®)

A. Criteria For Initial Approval

- i. Individual has a diagnosis of acromegaly; **AND**
- ii. Diagnosis of acromegaly has been confirmed by, or in consultation with, a board-certified endocrinologist who has reviewed and verified the test results (including, but not limited to: Insulin-like Growth Factor 1 levels; Oral Glucose Tolerance Test with associated Growth Hormone (GH) levels) that are indicative of a positive test; **AND**
- iii. Individual has had an inadequate response to any of the following:
 - A. Surgical resection; **OR**
 - B. Pituitary irradiation; **OR**
 - C. Bromocriptine mesylate at maximally tolerated doses;

OR
- iv. Surgery and/or radiotherapy is not an option;

OR

- v. Individual has a diagnosis of carcinoid tumors and is using for any of the following:
 - A. Metastatic carcinoid tumor to suppress or inhibit severe diarrhea and flushing episodes associated with the disease;

OR
- vi. Individual has a diagnosis of neuroendocrine and adrenal tumors and is using for any of the following:
 - A. For the management of unresectable locoregional disease or distant metastasis (NCCN 2A); **OR**
 - B. For the treatment of profuse watery diarrhea associated with VIPomas; **OR**

Medical Policy

Healthcare Services Department

Policy Name	Policy Number	Scope
Octreotide Agents (Sandostatin®, Sandostatin LAR®)	MP-RX-FP-65-23	<input checked="" type="checkbox"/> MMM MA <input checked="" type="checkbox"/> MMM Multihealth

C. Prophylactic treatment prior to surgery for gastrinoma (AHFS);

OR

- vii. Individual is using for bleeding Gastroesophagel (GE) varices and the following criteria are met:

A. Gastroesophageal varices are associated with liver disease (Banares 2002, Corley 2001);

AND

B. Octreotide acetate is used in combination with endoscopic therapy or alone if endoscopic therapy is not immediately available (Garcia-Tsao 2007);

OR

- viii. Individual is using for malignant bowel obstruction to manage gastrointestinal symptoms (e.g. nausea, pain, or vomiting) (AHFS);

OR

- ix. Individual is using for thymic carcinoma or thymoma with or without prednisone (NCCN 2A);

OR

- x. Individual is requesting Sandostatin LAR for meningiomas in central nervous system cancers (NCCN 2A); **AND**
- xi. Individual has surgically inaccessible recurrent or progressive disease when radiation is not possible; **AND**
- xii. Individual is using in combination with everolimus;

OR

- xiii. Individual is requesting Bynfezia or Sandostatin for rapid relief of symptoms or for breakthrough symptoms in individuals taking long-acting octreotide acetate or lanreotide when any of the criteria are met for the above uses (NCCN 2A).

B. Criteria For Continuation of Therapy

- i. MMM considers continuation of octreotide agents (Sandostatin®, Sandostatin LAR®) therapy medically necessary in members requesting reauthorization for an indication listed in Section A above (Criteria for Initial Approval) if the following information is provided:
 - a. Documentation of the patient's response to treatment showing no severe or life-threatening infusion reaction

C. Authorization Duration

- i. Initial Approval Duration: Up to 12 months
- ii. Reauthorization Approval Duration: Up to 12 months

D. Conditions Not Covered

Any other use is considered experimental, investigational, or unproven, including the following (this list may not be all inclusive)

Medical Policy

Healthcare Services Department

Policy Name	Policy Number	Scope
Octreotide Agents (Sandostatin®, Sandostatin LAR®)	MP-RX-FP-65-23	<input checked="" type="checkbox"/> MMM MA <input checked="" type="checkbox"/> MMM Multihealth

- i. Individual is using for the treatment of chylothorax; **OR**
- ii. Individual is using for the treatment of diarrhea associated with acquired immunodeficiency disease; **OR**
- iii. Individual is using for the treatment of gastrointestinal diseases (e.g. bleeding from vascular malformations, gastroparesis, pancreatitis, prevention of postoperative complications following pancreatic surgery, short bowel syndrome, or upper GI bleeding); **OR**
- iv. Individual is using for the treatment of Graves' ophthalmopathy; **OR**
- v. Individual is using for the treatment of hypothalamic obesity; **OR**
- vi. Individual is using for the treatment of other carcinomas (e.g. advanced breast cancer, hepatocellular cancer, or prostate cancer); **OR**
- vii. Individual is using for the treatment of polycystic kidney disease; **OR**
- viii. When the above criteria are not met and for all other indications.

Limits or Restrictions

A. Therapeutic Alternatives

The list below includes preferred alternative therapies recommended in the approval criteria and may be subject to prior authorization.

N/A

B. Quantity Limitations

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines. The chart below includes dosing recommendations as per the FDA-approved prescribing information.

Use	Recommended Dosing Schedule
Sandostatin® Injection	
Acromegaly	<ul style="list-style-type: none"> Initial dosage: 50 mcg subcutaneously three times daily for the first two weeks of therapy Maintenance dosage: 100 – 500 mcg three times daily.
Carcinoid Tumors	<ul style="list-style-type: none"> Initial dosage: 100 – 600 mcg subcutaneously daily in two to four divided doses during the initial 2 weeks of therapy. In the clinical studies, the median daily maintenance dosage was approximately 450 mcg but clinical and biochemical benefits were obtained in some patients with as little as 50 mcg, while others required doses up to 1500 mcg/day. However, experience with doses above 750 mcg/day is limited.

Medical Policy

Healthcare Services Department

Policy Name	Policy Number	Scope
Octreotide Agents (Sandostatin®, Sandostatin LAR®)	MP-RX-FP-65-23	<input checked="" type="checkbox"/> MMM MA <input checked="" type="checkbox"/> MMM Multihealth

VIPomas	<ul style="list-style-type: none"> Initial dosage: 200 – 300 mcg daily in two to four divided doses during the initial 2 weeks of therapy. Dosage may be adjusted to achieve a therapeutic response, but usually doses above 450 mcg/day are not required.
Sandostatin® LAR Injection	
Acromegaly	<ul style="list-style-type: none"> <i>Patients not currently receiving Sandostatin injection sc:</i> 50 mcg three times daily Sandostatin Injection subcutaneously for 2 weeks followed by SANDOSTATIN LAR DEPOT 20 mg intragluteally every 4 weeks for 3 months. <i>Patients currently receiving Sandostatin injection sc:</i> 20 mg every 4 weeks for 3 months.
Carcinoid Tumors and VIPomas	<ul style="list-style-type: none"> <i>Patients not currently receiving Sandostatin injection sc:</i> Sandostatin Injection subcutaneously 100 to 600 mcg/day in 2-4 divided doses for 2 weeks followed by SANDOSTATIN LAR DEPOT 20 mg every 4 weeks for 2 months. <i>Patients currently receiving Sandostatin injection sc:</i> 20 mg every 4 weeks for 2 months.
Exceptions for Sandostatin® LAR	
<ul style="list-style-type: none"> Renal Impairment, Patients on Dialysis: 10 mg every 4 weeks. Hepatic Impairment, Patients with Cirrhosis: 10 mg every 4 weeks. Quantity limits: <ul style="list-style-type: none"> Sandostatin LAR (octreotide) Depot Kit 20 mg: 2 kits per 28 days Sandostatin LAR (octreotide) Depot Kit 10 mg, 30 mg: 1 kit per 28 days Bynfezia (octreotide acetate) Pen 2,500mcg/ml (7,000mcg/2.8mL): 1 pen per 14 days 	

Medical Policy

Healthcare Services Department

Policy Name	Policy Number	Scope
Octreotide Agents (Sandostatin®, Sandostatin LAR®)	MP-RX-FP-65-23	<input checked="" type="checkbox"/> MMM MA <input checked="" type="checkbox"/> MMM Multihealth

Reference Information

1. DailyMed. Package inserts. U.S. National Library of Medicine, National Institutes of Health website. <http://dailymed.nlm.nih.gov/dailymed/about.cfm>. Accessed: June 25, 2024.
2. DrugPoints® System [electronic version]. Truven Health Analytics, Greenwood Village, CO. Updated periodically.
3. Lexi-Comp ONLINE™ with AHFS™, Hudson, Ohio: Lexi-Comp, Inc. Updated periodically.
4. NCCN Drugs & Biologics Compendium (NCCN Compendium™) © 2024 National Comprehensive Cancer Network, Inc. Available at: NCCN.org. Updated periodically.
5. Archimandritis A, Tsirantonaki M, Tryphonos M, et al. Ranitidine versus ranitidine plus octreotide in the treatment of acute nonvariceal upper gastrointestinal bleeding: a prospective randomised study. *Curr Med Res Opin.* 2000; 16(3):178-183.
6. Bajetta E, Procopio G, Ferrari L, et al. A randomized, multicenter prospective trial assessing long-acting release octreotide pamoate plus tamoxifen as a first line therapy for advanced breast carcinoma. *Cancer.* 2002; 94(2):299-304.
7. Banares R, Albillos A, Rincon D, et al. Endoscopic treatment versus endoscopic plus pharmacologic treatment for acute variceal bleeding: a meta-analysis. *Hepatology.* 2002; 35(3):609-615.
8. Berger J, Lester P, Rodrigues L. Medical Therapy of Malignant Bowel Obstruction With Octreotide, Dexamethasone, and Metoclopramide. *Am J Hosp Palliat Care.* 2016 May;33(4):407-410. Accessed: July 5, 2022.
9. Broder MS, Chang E, Cherepanov D, Neary MP, Ludlam WH. INCIDENCE AND PREVALENCE OF ACROMEGALY IN THE UNITED STATES: A CLAIMSBASED ANALYSIS. *Endocr Pract.* 2016; 22: 1327-1335.
10. Brown C, Subramanian V, Wilcox CM, Peter S. Somatostatin analogues in the treatment of recurrent bleeding from gastrointestinal vascular malformations: an overview and systematic review of prospective observational studies. *Dig Dis Sci.* 2010; 55(8):2129- 2134.
11. Caroli A, Perico N, Perna A, et al. Effect of longacting somatostatin analogue on kidney and cyst growth in autosomal dominant polycystic kidney disease (ALADIN): a randomised, placebo-controlled, multicentre trial. *Lancet.* 2013; 382(9903):1485-1495.
12. Chapman JA, Costantino JP, Dong B, et al. Octreotide LAR and tamoxifen versus tamoxifen in phase III randomize early breast cancer trials: NCIC CTG MA.14 and NSABP B-29. *Breast Cancer Res Treat.* 2015; 153(2):353-360.
13. Corley DA, Cello JP, Adkisson W, et al. Octreotide for acute esophageal variceal bleeding: a meta-analysis. *Gastroenterology.* 2001; 120(4):946-954.
14. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.: 2022. URL: <http://www.clinicalpharmacology.com>. Updated periodically.
15. DailyMed. Package inserts. U.S. National Library of Medicine, National Institutes of Health website. <http://dailymed.nlm.nih.gov/dailymed/about.cfm>. Accessed: July 5, 2022.
16. Das A, Shah PS. Octreotide for the treatment of chylothorax in neonates. *Cochrane Database Syst Rev.* 2010;(9):CD006388.
17. DrugPoints® System [electronic version]. Truven Health Analytics, Greenwood Village, CO. Updated periodically.

Medical Policy

Healthcare Services Department

Policy Name	Policy Number	Scope
Octreotide Agents (Sandostatin®, Sandostatin LAR®)	MP-RX-FP-65-23	<input checked="" type="checkbox"/> MMM MA <input checked="" type="checkbox"/> MMM Multihealth

18. Edmunds MC, Chen JD, Soykan I, et al. Effect of octreotide on gastric and small bowel motility in patients with gastroparesis. *Aliment Pharmacol Ther.* 1998; 12(2):167-174.
19. Friedlander TW, Weinberg VK, Small EJ, et al. Effect of the somatostatin analog octreotide acetate on circulating insulin-like growth factor-1 and related peptides in patients with non-metastatic castration-resistant prostate cancer: results of a phase II study. *Urol Oncol.* 2012; 30(4):408-414.
20. Fujita T, Daiko H. Efficacy and predictor of octreotide treatment for postoperative chylothorax after thoracic esophagectomy. *World J Surg.* 2014; 38(8):2039-2045.
21. Garcia-Tsao G, Sanyal AJ, Grace ND, et al. Prevention and management of gastroesophageal varices and variceal hemorrhage in cirrhosis. *Am J Gastroenterol.* 2007; 102(9):2086-2102.
22. Hogan MC, Masyuk TV, Page LJ, et al. Randomized clinical trial of long-acting somatostatin for autosomal dominant polycystic kidney and liver disease. *J Am Soc Nephrol.* 2010; 21(6):1052-1061
23. Ismail NA, Gordon J, Dunning J. The use of octreotide in the treatment of chylothorax following cardiothoracic surgery. *Interact Cardiovasc Thorac Surg.* 2015; 20(6):848-854.
24. Jia WD, Zhang CH, Xu GL, et al. Octreotide therapy for hepatocellular carcinoma: a systemic review of the evidence from randomized controlled trials. *Hepatogastroenterology.* 2010; 57(98):292-299.
25. unquera F, Saperas E, Videla S, et al. Long-term efficacy of octreotide in the prevention of recurrent bleeding from gastrointestinal angiodysplasia. *Am J Gastroenterol.* 2007; 102(2):254-260.
26. Lexi-Comp ONLINE™ with AHFS™, Hudson, Ohio: Lexi-Comp, Inc.; 2022; Updated periodically.
27. Loyaga-Rendon RY, Hashim T, Tallaj JA, et al. Octreotide in the management of recurrent gastrointestinal bleed in patients supported by continuous flow left ventricular assist devices. *ASAIO J.* 2015; 61(1):107-109.
28. Lustig RH, Hinds PS, Ringwald-Smith K, et al. Octreotide therapy of pediatric hypothalamic obesity: a double-blind, placebocontrolled trial. *J Clin Endocrinol Metab.* 2003; 88(6):2586-2592.
29. Nehra V, Camilleri M, Burton D, et al. An open trial of octreotide long-acting release in the management of short bowel syndrome. *Am J Gastroenterol.* 2001; 96(5):1494-1498.
30. NCCN Drugs & Biologics Compendium (NCCN Compendium™) © 2020 National Comprehensive Cancer Network, Inc. Available at: NCCN.org. Updated periodically.
31. Nikou GC, Toumpanakis C, Katsiari C, et al. Treatment of small intestinal disease in systemic sclerosis with octreotide: a prospective study in seven patients. *J Clin Rheumatol.* 2007; 13(3):119-123.
32. Panel on opportunistic infections in HIV-infected adults and adolescents. Guidelines for the prevention and treatment of opportunistic infections in HIV-infected adults and adolescents: recommendations from the Centers for Disease Control and Prevention, the National Institutes of Health, and the HIV Medicine Association of the Infectious Diseases Society of America. Last updated May 29, 2018. Available at: http://aidsinfo.nih.gov/contentfiles/lvguidelines/adult_oi.pdf.
33. Ruggerenti P, Remuzzi A, Ondei P, et al. Safety and efficacy of long-acting somatostatin treatment in autosomal-dominant polycystic kidney disease. *Kidney Int.* 2005; 68(1):206-216.

Medical Policy

Healthcare Services Department

Policy Name	Policy Number	Scope
Octreotide Agents (Sandostatin®, Sandostatin LAR®)	MP-RX-FP-65-23	<input checked="" type="checkbox"/> MMM MA <input checked="" type="checkbox"/> MMM Multihealth

34. Soudah HC, Hasler WL, Owyang C. Effect of octreotide on intestinal motility and bacterial overgrowth in scleroderma. N Engl J Med. 1991; 325(21):1461-1467.
35. Stan MN, Garrity JA, Bradley EA, et al. Randomized, double-blind, placebo-controlled trial of long-acting release octreotide for treatment of Graves' ophthalmopathy. J Clin Endocrinol Metab. 2006; 91(12):4817-4824.
36. Swanson MS, Hudson RL, Bhandari N, et al. Use of octreotide for the management of chyle fistula following neck dissection. JAMA Otolaryngol Head Neck Surg. 2015; 141(8):723-727.
37. Szilagyi A, Ghali MP. Pharmacological therapy of vascular malformations of the gastrointestinal tract. Can J Gastroenterol. 2006; 20(3):171-178.
38. Testoni D, Hornik CP, Neely ML, et al. Best Pharmaceuticals for Children Act - Pediatric Trials Network Administrative Core Committee. Safety of octreotide in hospitalized infants. Early Hum Dev. 2015; 91(7):387-392.
39. Verne GN, Eaker EY, Hardy E, Sninsky CA. Effect of octreotide and erythromycin on idiopathic and scleroderma-associated intestinal pseudoobstruction. Dig Dis Sci. 1995; 40(9):1892-1901.
40. Xu W, Zhou YF, Xia SH. Octreotide for primary moderate to severe acute pancreatitis: a meta-analysis. Hepatogastroenterology. 2013; 60(126):1504-1508

Federal and state laws or requirements, contract language, and Plan utilization management programs or policies may take precedence over the application of this clinical criteria.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

© CPT Only – American Medical Association

Medical Policy

Healthcare Services Department

Policy Name	Policy Number	Scope
Octreotide Agents (Sandostatin®, Sandostatin LAR®)	MP-RX-FP-65-23	<input checked="" type="checkbox"/> MMM MA <input checked="" type="checkbox"/> MMM Multihealth

Policy History

Revision Type	Summary of Changes	P&T Approval Date	UM/CMPC Approval Date
Annual Review 9/22/2025	Update octreotide agents to include FDA approved Bynfezia Pen (previously discontinued in 2021) within PA criteria and quantity limits. Also clarify use of Sandostatin for rapid relief of symptoms when using with long-acting octreotide or lanreotide. Wording and formatting updates. Coding Reviewed: Added HCPCS NOC C9399 and J3490 for Bynfezia Pen. Removed ICD-10-CM code range C7B.00-C7B.8 to add C7B.00-C7B.09 and C7B.8. Removed E34.0 and added E34.00-E34.09. Added C74.00-C74.92 and I85.01.	10/7/2025	10/17/2025
Annual Review 10/11/2024	<ul style="list-style-type: none"> Update prior 2A recommendation from NCCN for use with Sandostatin LAR product in CNS Meningiomas. Wording and formatting. Coding Reviewed: Updated HCPCS coding description for J2353 to clarify product is Sandostatin LAR depot; updated coding description for J2354 to clarify product is Sandostatin. Added ICD-10-CM D32.0-D32.9. Add NCCN 2A criteria for combination use with everolimus for CNS meningiomas. Wording and formatting updates. Removed obsolete agent, Bynfezia. Wording and formatting changes. Coding Reviewed: Removed Bynfezia from HCPCS J2354. 	11/18/2024	12/17/2024
Policy Inception	Elevance Health's Medical Policy adoption.	N/A	11/30/2023